

**Research Article**

## Physico-chemical characterization and Phytochemical Analysis of Leaves of Rejuvenative Herb, *Emblica officinalis*

Reni Nigam\*, Chanda Tiwari, Ranu Soni

Department of Biotechnology, Faculty of Life Sciences and Technology, AKS University, Satna (MP) 485001

<sup>1</sup>Corresponding Author E-mail: [reninigam83@gmail.com](mailto:reninigam83@gmail.com)

Received on: 18.12.20; Revised on: 19.01.21; Accepted on: 25.01.21

### Abstract

Towards authentication and quality assurance of medicinal herb, physicochemical and preliminary phytochemical studies of *Emblica officinalis* leaves were carried out. Physicochemical analysis revealed values for moisture contents, total ash, water soluble ash, acid insoluble ash, and water and alcohol extractives. Preliminary phytochemical screening revealed the presence of phytoconstituents viz., alkaloids, phenols, saponins, carbohydrates, proteins, in aqueous and hydroalcoholic extracts of *Emblica officinalis* leaves whereas flavonoids was present only in hydroalcoholic extract. Results showed that hydroalcoholic extract contain much more chemical constituents compare to aqueous extract. It was suggested that hydroalcoholic extract is the best solvent for extraction. Total ash (3.1%) was low implying that the plant leaves have low inorganic components. Water soluble ash and acid insoluble ash were found to be 1.7% and 1.3% respectively and water-soluble extractive value was low 28.8% compared to alcohol soluble extractive value 59.2%. Information obtained from present studies can be used in laying down identification and standardization of this plant as herbal remedy and also towards monograph development.

**Keywords:** Aqueous extract, *Emblica officinalis*, Physico-chemical, Phytochemical screening, Hydroalcoholic extract

### Introduction

Mother nature has gifted many medicinal plants to encourage healthy life. Numerous medicinal plants are used for herbal preparations in Ayurveda. *Phyllanthus emblica* Linn. Or *Emblica officinalis* Gaertn. (Amla), commonly known as Indian gooseberry belongs to the family Phyllanthaceae widely distributed in tropical and subtropical area

of India, China, Malaysia, Indonesia (Liu et. al., 2008). Amla is an important horticulture crop of arid zone. Owing to its multiple health benefits, it may be called as "King of Rasayana" [rejuvenation]. All parts of the plant are used in various Ayurvedic herbal preparations, including the fruit, seed, leaves, root, bark and flower. The medicinal preparation from amla, have high potency in curing many diseases and promotion of health (Rajeshkumar et al., 2001) and used widely in combination with

other two fruit-bearing plant species, chebulic and belleric as triphala. It is key ingredient in the Chyavanaprasha (ayurvedic recipe) (Singh et al., 2011). In India, fruits of *Emblica officinalis* are the most common ingredients of almost all Ayurvedic preparations like Lehya, Choorna etc. (Raghu and Ravindra, 2010). It contains polyphenols, flavonoids, tannins, pectin, gallic acid, emblicanin, punigluconin and other bioactive compounds (Zhang et al., 2003, Prashant, 2014). It has many therapeutic applications such as hepatoprotective, Immunostimulants, antimicrobial, antioxidants, anticancer, antipyretics, cytoprotective, analgesic, antitussive, anti-inflammatory, gastroprotective and used in eye disorder (Khan, 2009). It is found to be effective for the reversal of dyslipidemia and intima-media thickening and plaque formation in the aorta in hypercholesterolaemic rabbits (Antony, 2006). Antioxidants present in its juice in the form of polyphenols and vitamin C help in maintaining healthy cholesterol level thus provide a cardio-protective effect (Pathak et al., 2003). The antibacterial activity of aqueous crude extracts of *Emblica officinalis* against human bacterial pathogens was reported (Kanthimathi, 2013, Rawat, 2015). Many chemical constituents present in its different parts like leaves-contain chebulic acid, gallic acid, chebulinic acid, ellagic acid, amlic acid, phyllantin, alkanoids, seeds-contain A phosphalides, fixed oil and a small quantity of essential oil barks-contain proanthocyanin, leukodelphinidin and tannin and roots-contain luprol and ellegic acid (Khan, 2009). A variety of phytochemicals such as tannins, flavonoids, terpenoids and alkaloids were reported to have several pharmacological properties such as antioxidant, anticancer, antitumor, antigenotoxic and anticarcinogenic effects. It was considered to be a safe herbal medicine without any adverse effects. Hence, the present study deals with the screening of phytochemical constituents present in the aqueous and hydroalcoholic extracts of *Emblica officinalis* leaves and their physicochemical characterization.

## Materials and Methods

### Plant material

The leaves of plant were collected from local area of Satna district and identification of the plant

sample was done. The shade dried leaves were cut into small pieces and made into coarsely powder using mechanical grinder and preserved in air tight container (Harborne, 2007).

### Plant Extracts Preparation

For preliminary phytochemical screening dried and coarsely powered leaves (50 g/250 mL) was extracted with solvents (water and hydroalcoholic) in a Soxhlet extractor by continuous hot percolation process. The extracts were evaporated for dryness in water bath and used for phytochemical determination (Harborne, 2007).

### Phytochemical Screening

The extracts of *Emblica officinalis* leaves were analyzed for the presence of various phytoconstituents by procedure given in Mukherjee (2007), Khandelwal (2005), Kokate (2011).

### Test for Alkaloids

Plant extracts were dissolved individually in dilute hydrochloric acid (dil. HCl) and filtered. Mayer's test: In this test procedure, filtrates were treated with Mayer's reagent (Potassium mercuric iodide) after few minutes yellow colored precipitate appeared in test tube indicates the presence of alkaloids in plant sample. Hager's test: In Hager's test, extract was treated with Hager's reagent (saturated picric acid solution) and presence of the alkaloids in sample confirmed by the formation of yellow colored precipitate in test tube.

### Test for Carbohydrates

Extracts were dissolved individually in 5 ml distilled water and filtrated. The filtrates were used to test for the presence of carbohydrates. In Fehling test, firstly plant extract was taken in test tube and hydrolyzed with dilute HCl, neutralized with alkali and heated with Fehling's A & B solution, red color precipitate appeared in test tube indicates the presence of reducing sugar in plant sample.

### Test for Saponins

For determination of saponins in plant sample, extracts were diluted with distilled water up to 20 ml in test tube and was shaken in a graduated cylinder for 15 minutes after some time 1 cm layer of foam form in test tube, indicates the presence of saponins in plant sample. In another foam test, 0.5gm of plant extract was shaken with 2 ml of distilled water. If foam produced and persists for 10

minutes, it indicates the presence of saponins in plant sample.

### Test for Phenols

In this test, plant extracts were treated with 3 to 4 drops of ferric chloride solution; bluish black color form in the test tube confirmed the presence of phenols.

### Test for Flavonoids

In alkaline reagent test, firstly, take 2 ml of test solution in test tube, add few drops of sodium hydroxide solution, intense yellow color was formed which turns to colorless on addition of few drops of dilute acid indicates presence of flavonoids. In lead acetate test, firstly, take 2 ml of plant extract solution subsequently, add few drops of lead acetate solution in test tube, yellow colour precipitate appeared in test tube indicates the presence of flavonoids in plant sample.

### Test for Proteins

Xanthoproteic test: In this test procedure, 5 ml of test solution taken in test tube, and add 1 ml of concentrated nitric acid and boil, yellow color precipitate was formed in test tube. After cooling this test solution, add 40% sodium hydroxide solution, orange color precipitate formed confirmed presence of protein in plant sample.

### Physico-Chemical Analysis

Physico-chemical analysis of plant sample was carried out according to the standard method of Indian Pharmacopoeia, (2010) and Tripathi and Sikarwar, (2014).

### Determination of Moisture (Loss of Drying)

Two gram of powdered plant material was weighted and dried in an oven at (100°C) until two consecutive weighing do not differ by more than 2gm then, loss of weight of plant sample calculated in terms of percentage.

### Ash value

About three gram of powdered plant sample was weighed into a tarred silica crucible and incinerated at 450°C in a muffle furnace until free from carbon. Subsequently, the crucible was cooled and weighed and percentage of total ash was calculated with compared to air dried sample.

### Water Soluble Ash

1.5 gm ash obtained from the total ash was boiled with 25 ml of distilled water for (8 minutes) and

filtered through an ash less filter paper and wash two times with hot water. The filter paper was transferred into a tarred silica crucible, incinerated at 450°C in a muffle furnace until free from carbon. The crucible was cooled and weighed and percentage of water-soluble ash calculated with reference to air-dried substance.

### Acid Insoluble Ash

1.5 gm ash obtained from the total ash was boiled with 25 ml of 2N HCl (for 8 minutes). Subsequently, filter through an ash less filter paper and wash filter paper two times with hot water. The filter paper was transferred into a tarred silica crucible, incinerated at 450°C in a muffle furnace until free from carbon. The crucible was cooled and weighed and percentage of acid insoluble ash was calculated with reference to air-dried substance.

### Determination of Alcohol and Water-Soluble Extractive

About 5 g of air-dried plant material was macerate with 100 ml of the solvents (ethanol and water) for 6 hours, shake frequently then allowed to stand for 18 hours in room temperature. Then, filtered rapidly and transferred 25 ml of the filtrate to a tarred flat-bottomed petri-dish and evaporate to dry on a water bath at 100°C for 6 hours, cool in desiccators for 30 minutes and weighed and calculated as milligram per ml of extractive values in terms of percentage.

## Results and Discussion

Medicinal plants are potential renewable natural resources and are generally considered to play a beneficial role in human health care. The medicinal value of this plant lies in some chemical substances that produce a definite physiological action on the human body. The most important of these bioactive compounds are alkaloids, flavonoids, tannins and phenols. The results of present study showed that *Emblia officinalis* is rich in phytochemical and physicochemical contents as highlighted in table 1 and 2. Multifunctional activities of the plants against various diseases may be attributed due to the presence of various chemical constituents in the plant. Preliminary phytochemical revealed the presence of phytoconstituents viz., alkaloids, phenols, saponins, carbohydrates, proteins in aqueous and

hydroalcoholic extracts of leaves of *Emblica officinalis* whereas flavonoids was present only in hydroalcoholic extract. Similar observations were made by Khopde Priyadarsini, (2001) and Patil, (2013). The presence of flavonoids in *Emblica officinalis* is likely to be responsible for the free radical scavenging effects. The plant also contains phenolics which are powerful antioxidants. The presence of saponins protects plant from microbial pathogen. It showed that hydroalcoholic extract contain much more phytochemical constituents than aqueous extract of leaves of this plant. Another study of Prashant, (2014) also revealed the presence of phytochemicals like flavonoid, alkaloids and phenols in *Emblica officinalis* leaves extract.

The determination of physicochemical parameter is an important in determination of adulterants and improper handling of drugs. The percentage of alcohol soluble extractive value and water-soluble extractive value were also determined and results

are depicted in Table 2. The LOD results 10%, was an indication that the extract was least hygroscopic and the plant can be stored for a long period of time with less probability of microbial attack. Total ash of 3.1 % was low implying that the crude plant has low inorganic components. Water soluble ash and acid insoluble ash found to be 1.7% and 1.3% respectively. Acid insoluble ash is less than the water-soluble ash value. Water soluble extractive value is less 28.8% compared to alcohol soluble extractive value of 59.2% which shows that aqueous solvent not permeates the cells of leaves of *Emblica officinalis*. The results of percentage extractive yield for *Emblica officinalis* indicate that plant sample was highly soluble in alcohol than water solvents.

Table No. 1 Qualitative Phytochemical Screening of leaves extracts of *Emblica officinalis*

Chemical Constituents	Test/ Reagent	Results	
		Aqueous extract	Hydroalcoholic extract
Alkaloids	Hager's Test	+ ve	+ ve
	Wagner's Test	+ ve	+ ve
Carbohydrate	Fehling's Test	+ ve	+ ve
Phenols	Fecl <sub>3</sub> Test	- ve	+ ve
	Alkaline Test	- ve	+ ve
Flavonoids	Lead Acetate Test	- ve	+ ve
Saponins	Foam Test	+ ve	+ ve
Proteins	Xanthoproteic	+ ve	+ ve
Diterpenes	Copper Acetate Test	- ve	- ve

(+) **Ve**: Indicates the presence of phytochemicals, (-) **Ve**: Indicates the absence of phytochemicals

Table No. 2 Physico-Chemical characteristics of leaves of *Emblica officinalis*

Parameter	Results (%) W/W
<b>Ash Value</b>	
Total Ash	3.1%
Acid Insoluble Ash Value	1.3%
Water Soluble Ash Value	1.7%
<b>Extractive Value</b>	
Ethanol Soluble Extractive Value	59.2%
Water Soluble Extractive Value	28.8%
<b>Loss on Drying</b>	10%

## CONCLUSION

For healthy look and fitness, present generation is using supplementary vitamin pills to sustain and boost their system according to medical consultation, they could even prove to be fatal for life if they don't choose the nature's products. Despite various available allopathic formulations, the plant-based compounds are beneficial for various diseases, the present study was undertaken to reveal the physico-chemical and phytochemical characterization which could subsequently be exploited as cost-effective measures for health of well-being.

## ACKNOWLEDGEMENT

The authors are grateful to Mr. Anant Soni, Pro-Chancellor, AKS University, Satna and Dr. Kamlesh Choure, Head, Department of Biotechnology, AKS University, Satna, India for providing the necessary laboratory facilities and for his valuable guidance.

## CONFLICT OF INTEREST

The author declares no conflict of interest.

## REFERENCE

- Anonymous, Laboratory guideline for the analysis of Ayurveda and Siddha formulations, (2010). Central Council for Research in Ayurveda and Siddha. Department of Ayush, Ministry of Health and Family Welfare, Government of India, New Delhi.
- Antony, B., Merina, B., Sheeba, V., Mukkadan, J. (2006). Effect of standardized Amla extract on atherosclerosis and dyslipidemia. *Ind J Pharm Sci*, 68(4), 437–441.
- Arora, A., Kumar, I., Sen, R., and Singh, J. (2011). *Emblica officinalis* (Amla): Physicochemical and fatty acid analysis from arid Zone of Rajasthan. *International Journal of Basic and Applied Chemical Sciences*, 72, 281-286.
- Harborne, J.B. (2007). *Phytochemical methods, A guide to modern techniques of plant analysis*, 3rd ed. Chapman and Hall: London, 16-32.
- Kanthimathi, M., and Soranam, R. (2013). Antibacterial effects of *Emblica officinalis* and *Phyllanthus niruri* crude extracts against bacterial pathogens. *International Journal of Pharmaceutical and Clinical Science*, 3(3), 20-23.
- Khan, K.H. (2009). Roles of *Emblica officinalis* in Medicine- A Review. *Botany Research International Publications*, 2(4), 218-228.

- Khandelwal, K.R. (2005). Practical pharmacognosy, technique and experiments, 23rd Ed. Nirali Prakashan.
- Khopde, S.M., Indira Priyadarsini, K., Mohan, H., Gawandi, V.B., Satav, J.G., Yakhmi, J.V., Banavaliker, M.M., Biyani, M.K., Mittal, J.P. (2001). Characterizing the antioxidant activity of amla (*Phyllanthus emblica*) extract. *Current science*, 81(2), 185-190.
- Kokate, C.K. (1994). Practical Pharmacognosy, 4th Edn. Vallabh Prakashan, 112-120.
10. Kumar, G.S., Nayakaa, H., Shylaja, D.M., Salimatha, P.V. (2006). Free and bound phenolic antioxidants in Amla (*Embolica officinalis*) and turmeric (*Curcuma longa*). *J. Food Comp*, 19, 446-452.
- Liu, X., Cui, C., ZHAO, M., Wang, J., LUO, W., YANG, B., Jiang, Y. (2008). Identification of phenolics in the fruit of *Embolica* (*Phyllanthus emblica* L.) and their activities. *Food Chem*, 109, 909-915.
- Meena, A.K., and Singh, S. (2010). Evaluation of physicochemical and preliminary phytochemical studies on the fruit of *Embolica officinalis* Gaertn. *Asian Journal of Pharmaceutical and Clinical Research*, 3(3), 242-243.
- Mukherjee, P.K. (2007). Quality control of herbal drugs, 2nd Ed. Business Horizons.
- Pathak, R.K. (2003). Status report on genetic resources of Indian gooseberry-Aonla (*Embolica officinalis* Gaertn.) in south and south East Asia. IPGRI Office for South Asia National Agriculture Science Centre (NASC).
- Patil, S.G., Deshmukh, A.A., Padol, A.R., Jagadale, A.A. (2013). Phytochemical characterization and estimation of percent extractability of *Embolica officinalis* fruit extract. *International Journal of Natural Products Research*, 2(1), 20-24.
- Prashanth, G.K., Krishnaiah, G.M. (2014). Phytochemical screening and GC-MS analysis of the leaves of *Pongamia Pinnata* Linn. *International Journal of Innovative Research in Science*, 3(11), 17329-17333.
- Raghu, H.S., and Ravindra, P. (2010). Antimicrobial activity and phytochemical study of *Phyllanthus emblica* linn. *International Journal of Pharmaceutical Studies and Research*, 1(1), 30-33.
- Rajeshkumar, N.V., Therese, M., Kuttan, R. (2001). *Embolica officinalis* fruit effect protection against experimental gastric user in rats. *Pharma. Biol*, 39, 375-380.
- Rawat, S. (2015). Antimicrobial activity of Neem, Tulsi, Henna and Amla against pathogenic bacteria. *Journal of Chemical and Pharmaceutical Research*, 7(4), 1056-1059.
- Singh, E., Sharma, S., Pareek, A. (2011). Phytochemistry, traditional uses and cancer chemopreventive activity of Amla (*Phyllanthus emblica*): The Sustainer. *Journal of Applied Pharmaceutical Science*, 02(01), 176-183.
- Thomas, R.E., and Kamat, S.D. (2014). Microwave and ultrasonication assistant extraction of phytochemicals from *B. Monnieri* and study of its antioxidant activity. *International journal of pharma and bio sciences*, 5(1), 66-69.
- Tripathi, M.K and Sikarwar, R.L.S. (2014). Pharmacognostic standardization of stem bark of *Erythrina variegata* L. *The Journal of Indian Botanical Society*, 93(3&4), 248-253.
- Yadav, V., Duvey, B., Sharma, S., Devi, B. (2014). Amla (*Embolica officinalis*)- Medicinal Food and Pharmacological activity amla (*Embolica officinalis*)- Medicinal Food and pharmacological Activity. *International journal of pharmaceutical and chemical sciences*, 3(3), 616-619.
- Yokozawa, T., Kim, H.Y., Kim, H.J., Okubo, T., Chu, D.C., Juneja, L.R. (2007). Amla (*Embolica officinalis* Gaertn.) prevents dyslipidaemia and oxidative stress in the aging process. *British J. Nutri*, 97(6), 1187-1195.
- Zhang, L.Z., ZHAO, W.H., GUO, Y.J., TU, G.Z., Lin, S., XIN, L.G. (2003). Studies on chemical constituent in fruits of Tibetan medicine *Phyllanthus emblica*. *Zhongguo Zhong YaoZaZhi*, 28(10), 940-943.